

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
11 November 2004 (11.11.2004)

PCT

(10) International Publication Number
WO 2004/096804 A1

- (51) International Patent Classification⁷: **C07D 471/08**, A61K 31/4995, A61P 13/12
- (74) Agent: **HOFMANN, Dieter**; Therwilerstrasse 87, CH-4153 Reinach (CH).
- (21) International Application Number: **PCT/EP2004/004373**
- (22) International Filing Date: 26 April 2004 (26.04.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: **PCT/EP03/04393** 28 April 2003 (28.04.2003) EP
- (71) Applicant (for all designated States except US): **ACTELION PHARMACEUTICALS LTD** [CH/CH]; Gewerbestrasse 16, CH-4123 Allschwil (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **BEZENCON, Olivier** [CH/CH]; Dürrenmattweg 53, CH-4123 Allschwil (CH). **BUR, Daniel** [CH/CH]; Im Rosengarten 24, CH-4106 Therwil (CH). **FISCHLI, Walter** [CH/CH]; Obertorweg 64, CH-4123 Allschwil (CH). **REMEN, Lubos** [SK/CH]; Kurzellängeweg 28, CH-4123 Allschwil (CH). **RICHARD-BILDSTEIN, Sylvia** [FR/FR]; 12, rue des Beaux Prés, F-68440 Dietwiller (FR). **WELLER, Thomas** [CH/CH]; Hoezlistrasse 58, CH-4102 Binningen (CH).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DIAZABICYCLONONENE AND TETRAHYDROPYRIDINE DERIVATIVES AS RENIN INHIBITORS

(57) Abstract: The invention relates to novel 3,9-diazabicyclo[3.3.1]nonene derivatives, tetrahydropyridine derivatives, and related compounds and their use as active ingredients in the preparation of pharmaceutical compositions. The invention also concerns related aspects including processes for the preparation of the compounds, pharmaceutical compositions containing one or more of those compounds and especially their use as inhibitors of renin.

WO 2004/096804 A1

DIAZABICYCLONONENE AND
TETRAHYDROPYRIDINE DERIVATIVES AS RENIN INHIBITORS

5

The invention relates to novel compounds of the general formula I. The invention also concerns related aspects including processes for the preparation of the compounds, pharmaceutical compositions containing one or more compounds of formula I and especially their use as renin inhibitors in cardiovascular events and renal insufficiency. Furthermore, these compounds can be regarded as inhibitors of other aspartyl proteases and might therefore be useful as inhibitors of plasmepsins to treat malaria and as inhibitors of *Candida albicans* secreted aspartyl proteases to treat fungal infections.

15 In the renin-angiotensin system (RAS) the biologically active angiotensin II (Ang II) is generated by a two-step mechanism. The highly specific enzyme renin cleaves angiotensinogen to angiotensin I (Ang I), which is then further processed to Ang II by the less specific angiotensin-converting enzyme (ACE). Ang II is known to work on at least two receptor subtypes called AT₁ and AT₂. Whereas
20 AT₁ seems to transmit most of the known functions of Ang II, the role of AT₂ is still unknown.

Modulation of the RAS represents a major advance in the treatment of cardiovascular diseases. ACE inhibitors and AT₁ blockers have been accepted to
25 treat hypertension (Waeber B. *et al.*, "The renin-angiotensin system: role in experimental and human hypertension", in Berkenhager W. H., Reid J. L. (eds): *Hypertension*, Amsterdam, Elsevier Science Publishing Co, 1996, 489-519; Weber M. A., *Am. J. Hypertens.*, 1992, 5, 247S). In addition, ACE inhibitors are used for renal protection (Rosenberg M. E. *et al.*, *Kidney International*, 1994, 45,
30 403; Breyer J. A. *et al.*, *Kidney International*, 1994, 45, S156), in the prevention of congestive heart failure (Vaughan D. E. *et al.*, *Cardiovasc. Res.*, 1994, 28, 159;

Fouad-Tarazi F. *et al.*, *Am. J. Med.*, **1988**, *84 (Suppl. 3A)*, 83) and myocardial infarction (Pfeffer M. A. *et al.*, *N. Engl. J. Med.*, **1992**, 327, 669).

The rationale to develop renin inhibitors is the specificity of renin (Kleinert H. D.,
5 *Cardiovasc. Drugs*, **1995**, *9*, 645). The only substrate known for renin is
angiotensinogen, which can only be processed (under physiological conditions) by
renin. In contrast, ACE can also cleave bradykinin besides Ang I and can be by-
passed by chymase, a serine protease (Husain A., *J. Hypertens.*, **1993**, *11*, 1155).
In patients inhibition of ACE thus leads to bradykinin accumulation causing
10 cough (5-20%) and potentially life-threatening angioneurotic edema (0.1-0.2%)
(Israili Z. H. *et al.*, *Annals of Internal Medicine*, **1992**, *117*, 234). Chymase is not
inhibited by ACE inhibitors. Therefore, the formation of Ang II is still possible in
patients treated with ACE inhibitors. Blockade of the AT₁ receptor (e.g. by
losartan) on the other hand overexposes other AT-receptor subtypes to Ang II,
15 whose concentration is dramatically increased by the blockade of AT₁ receptors.
This may raise serious questions regarding the safety and efficacy profile of AT₁
receptor antagonists. In summary, renin inhibitors are not only expected to be
different from ACE inhibitors and AT₁ blockers with regard to safety, but more
importantly also with regard to their efficacy to block the RAS.

20

Only limited clinical experience (Azizi M. *et al.*, *J. Hypertens.*, **1994**, *12*, 419;
Neutel J. M. *et al.*, *Am. Heart*, **1991**, *122*, 1094) has been created with renin
inhibitors because of their insufficient oral activity due to their peptidomimetic
character (Kleinert H. D., *Cardiovasc. Drugs*, **1995**, *9*, 645). The clinical
25 development of several compounds has been stopped because of this problem
together with the high cost of goods. Only one compound containing four chiral
centers has entered clinical trials (Rahuel J. *et al.*, *Chem. Biol.*, **2000**, *7*, 493;
Mealy N. E., *Drugs of the Future*, **2001**, *26*, 1139). Thus, metabolically stable,
orally bioavailable and sufficiently soluble renin inhibitors that can be prepared on
30 a large scale are missing and sought. Recently, the first non-peptide renin
inhibitors were described which show high *in vitro* activity (Oefner C. *et al.*,
Chem. Biol., **1999**, *6*, 127; Patent Application WO97/09311; Märki H. P. *et al.*, *Il*

Farmaco, 2001, 56, 21). However, the development status of these compounds is not known.


The present invention relates to the identification of renin inhibitors of a non-peptidic nature and of low molecular weight. Orally active renin inhibitors of long duration of action which are active in indications beyond blood pressure regulation where the tissular renin-chymase system may be activated leading to pathophysiologically altered local functions such as renal, cardiac and vascular remodeling, atherosclerosis, and possibly restenosis are described.

10

The present invention describes non-peptidic renin inhibitors.

In particular, the present invention relates to novel compounds of the general formula I,

15



wherein

Z, Y, X and W represent independently a nitrogen atom or a -CH- group; at least two of the Z, Y, X and W represent a -CH- group;

V represents a bond; $-(CH_2)_r-$; $-A-(CH_2)_s-$; $-CH_2-A-(CH_2)_t-$; $-(CH_2)_s-A-$; $-(CH_2)_2-A-(CH_2)_u-$; $-A-(CH_2)_v-B-$; $-CH_2-CH_2-CH_2-A-CH_2-$; $-A-CH_2-CH_2-B-CH_2-$; $-CH_2-A-CH_2-CH_2-B-$; $-CH_2-CH_2-CH_2-A-CH_2-CH_2-$; $-CH_2-CH_2-CH_2-CH_2-A-CH_2-$; $-A-$

CH₂-CH₂-B-CH₂-CH₂-; -CH₂-A-CH₂-CH₂-B-CH₂-; -CH₂-A-CH₂-CH₂-CH₂-B-; or
-CH₂-CH₂-A-CH₂-CH₂-B-;

A and B independently represent -O-; -S-; -SO-; -SO₂-;

5

U represents aryl; heteroaryl;

T represents -CONR¹-; -(CH₂)_pOCO-; -(CH₂)_pN(R¹)CO-; -(CH₂)_pN(R¹)SO₂-; or
-COO-;

10

Q represents lower alkylene; lower alkenylene;

M represents hydrogen; cycloalkyl; aryl; heterocyclyl; heteroaryl;

15 L represents -R³; -COR³; -COOR³; -CONR²R³; -SO₂R³; -SO₂NR²R³;
-COCH(Aryl)₂;

R¹ represents hydrogen; lower alkyl; lower alkenyl; lower alkynyl; cycloalkyl;
aryl; cycloalkyl - lower alkyl;

20

R² and R^{2'} independently represent hydrogen; lower alkyl; lower alkenyl;
cycloalkyl; cycloalkyl - lower alkyl;

R³ represents hydrogen; lower alkyl; lower alkenyl; cycloalkyl; aryl; heteroaryl;
25 heterocyclyl; cycloalkyl - lower alkyl; aryl - lower alkyl; heteroaryl - lower alkyl;
heterocyclyl - lower alkyl; aryloxy - lower alkyl; heteroaryloxy - lower alkyl,
whereby these groups may be unsubstituted or mono-, di- or trisubstituted with
hydroxy, -OCOR², -COOR², lower alkoxy, cyano, -CONR²R^{2'}, -CO-morpholin-4-
yl, -CO-((4-loweralkyl)piperazin-1-yl), -NH(NH)NH₂, -NR⁴R^{4'} or lower alkyl,

30 with the proviso that a carbon atom is attached at the most to one heteroatom in
case this carbon atom is sp³-hybridized;

R^4 and $R^{4'}$ independently represent hydrogen; lower alkyl; cycloalkyl; cycloalkyl - lower alkyl; hydroxy - lower alkyl; $-COOR^2$; $-CONH_2$;

k is the integer 0 or 1;

5

m and n represent the integer 0 or 1, with the proviso that in case m represents the integer 1, n is the integer 0; in case n represents the integer 1, m is the integer 0; in case k represents the integer 0, n represents the integer 0;

- 10 p is the integer 1, 2, 3 or 4;
r is the integer 1, 2, 3, 4, 5, or 6;
s is the integer 1, 2, 3, 4, or 5;
t is the integer 1, 2, 3, or 4;
u is the integer 1, 2, or 3;
15 v is the integer 2, 3, or 4;

and optically pure enantiomers, mixtures of enantiomers such as racemates, diastereomers, mixtures of diastereomers, diastereomeric racemates, mixtures of diastereomeric racemates, and the meso-form; as well as pharmaceutically
20 acceptable salts, solvent complexes and morphological forms.

In the definitions of general formula I – if not otherwise stated – the term **lower alkyl**, alone or in combination with other groups, means saturated, straight and branched chain groups with one to seven carbon atoms, preferably one to four
25 carbon atoms that can be optionally substituted by halogens. Examples of lower alkyl groups are methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, pentyl, hexyl and heptyl. The methyl, ethyl and isopropyl groups are preferred.

30 The term **lower alkoxy** refers to a R-O-group, wherein R is a lower alkyl. Examples of lower alkoxy groups are methoxy, ethoxy, propoxy, iso-propoxy, isobutoxy, sec-butoxy and tert-butoxy.

The term **lower alkenyl**, alone or in combination with other groups, means straight and branched chain groups comprising an olefinic bond and consisting of two to seven carbon atoms, preferably two to four carbon atoms, that can be optionally substituted by halogens. Examples of lower alkenyl are vinyl, propenyl or butenyl.

The term **lower alkynyl**, alone or in combination with other groups, means straight and branched chain groups comprising a triple bond and consisting of two to seven carbon atoms, preferably two to four carbon atoms, that can be optionally substituted by halogens. Examples of lower alkynyl are ethynyl, propynyl or butynyl.

The term **lower alkylene**, alone or in combination with other groups, means straight and branched divalent chain groups with one to seven carbon atoms, preferably one to four carbon atoms, that can be optionally substituted by halogens. Examples of lower alkylene are ethylene, propylene or butylene.

The term **lower alkenylene**, alone or in combination with other groups, means straight and branched divalent chain groups comprising an olefinic bond and consisting of two to seven carbon atoms, preferably two to four carbon atoms, that can be optionally substituted by halogens. Examples of lower alkenylene are vinylene, propenylene and butenylene.

The term **lower alkylenedioxy**, refers to a lower alkylene substituted at each end by an oxygen atom. Examples of lower alkylenedioxy groups are preferably methylenedioxy and ethylenedioxy.

The term **lower alkyleneoxy** refers to a lower alkylene substituted at one end by an oxygen atom. Examples of lower alkyleneoxy groups are preferably methyleneoxy, ethyleneoxy and propyleneoxy.

The term **halogen** means fluorine, chlorine, bromine or iodine, preferably fluorine, chlorine and bromine.

The term **cycloalkyl** alone or in combination, means a saturated cyclic hydrocarbon ring system with 3 to 7 carbon atoms, e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl, which can be optionally mono- or multisubstituted by lower alkyl, lower alkenyl, lower alkenylene, lower alkoxy, lower alkyleneoxy, lower alkylenedioxy, hydroxy, halogen, $-\text{CF}_3$, $-\text{NR}^1\text{R}^{1'}$, $-\text{NR}^1\text{C}(\text{O})\text{R}^{1'}$, $-\text{NR}^1\text{S}(\text{O}_2)\text{R}^{1'}$, $-\text{C}(\text{O})\text{NR}^1\text{R}^{1'}$, lower alkylcarbonyl, $-\text{COOR}^1$, $-\text{SR}^1$, $-\text{SOR}^1$, $-\text{SO}_2\text{R}^1$, $-\text{SO}_2\text{NR}^1\text{R}^{1'}$ whereby $\text{R}^{1'}$ represents hydrogen; lower alkyl; lower alkenyl; lower alkynyl; cycloalkyl; aryl; cycloalkyl - lower alkyl. The cyclopropyl group is a preferred group.

The term **aryl**, alone or in combination, relates to the phenyl, the naphthyl or the indanyl group, preferably the phenyl group, which can be optionally mono- or multisubstituted by lower alkyl, lower alkenyl, lower alkynyl, lower alkenylene or lower alkylene forming with the aryl ring a five- or six-membered ring, lower alkoxy, lower alkylenedioxy, lower alkyleneoxy, hydroxy, hydroxy-lower alkyl, halogen, cyano, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{NR}^1\text{R}^{1'}$, $-\text{NR}^1\text{R}^{1'}$ - lower alkyl, $-\text{NR}^1\text{C}(\text{O})\text{R}^{1'}$, $-\text{NR}^1\text{S}(\text{O}_2)\text{R}^{1'}$, $-\text{C}(\text{O})\text{NR}^1\text{R}^{1'}$, $-\text{NO}_2$, lower alkylcarbonyl, $-\text{COOR}^1$, $-\text{SR}^1$, $-\text{SOR}^1$, $-\text{SO}_2\text{R}^1$, $-\text{SO}_2\text{NR}^1\text{R}^{1'}$, benzyloxy, whereby $\text{R}^{1'}$ has the meaning given above.

The term **aryloxy** refers to an Ar-O-group, wherein Ar is an aryl. An example of a lower aryloxy group is phenoxy.

25

The term **heterocyclyl**, alone or in combination, means saturated or unsaturated (but not aromatic) five-, six- or seven-membered rings containing one or two nitrogen, oxygen or sulfur atoms which may be the same or different and which rings can be optionally substituted with lower alkyl, hydroxy, lower alkoxy and halogen. The nitrogen atoms, if present, can be substituted by a $-\text{COOR}^2$ group. Examples of such rings are piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, dihydropyranyl, 1,4-dioxanyl, pyrrolidinyl,

tetrahydrofuranyl, dihydropyrrolyl, imidazolidinyl, dihydropyrazolyl, pyrazolidinyl, dihydroquinolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl.

The term **heteroaryl**, alone or in combination, means six-membered aromatic rings containing one to four nitrogen atoms; benzofused six-membered aromatic rings containing one to three nitrogen atoms; five-membered aromatic rings containing one oxygen, one nitrogen or one sulfur atom; benzofused five-membered aromatic rings containing one oxygen, one nitrogen or one sulfur atom; five-membered aromatic rings containing one oxygen and one nitrogen atom and benzofused derivatives thereof; five-membered aromatic rings containing a sulfur and a nitrogen or an oxygen atom and benzofused derivatives thereof; five-membered aromatic rings containing two nitrogen atoms and benzofused derivatives thereof; five-membered aromatic rings containing three nitrogen atoms and benzofused derivatives thereof, or a tetrazolyl ring. Examples of such ring systems are furanyl, thiophenyl, pyrrolyl, pyridinyl, pyrimidinyl, indolyl, quinolinyl, isoquinolinyl, imidazolyl, triazinyl, thiazinyl, thiazolyl, isothiazolyl, pyridazinyl, pyrazolyl, oxazolyl, isoxazolyl, coumarinyl, benzothiophenyl, quinazolinyl, quinoxalinyl. Such rings may be adequately substituted with lower alkyl, lower alkenyl, lower alkynyl, lower alkylene, lower alkenylene, lower alkylenedioxy, lower alkyleneoxy, hydroxy-lower alkyl, lower alkoxy, hydroxy, halogen, cyano, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{NR}^1\text{R}^{1'}$, $-\text{NR}^1\text{R}^{1'}$ - lower alkyl, $-\text{N}(\text{R}^1)\text{COR}^1$, $-\text{N}(\text{R}^1)\text{SO}_2\text{R}^1$, $-\text{CONR}^1\text{R}^{1'}$, $-\text{NO}_2$, lower alkylcarbonyl, $-\text{COOR}^1$, $-\text{SR}^1$, $-\text{SOR}^1$, $-\text{SO}_2\text{R}^1$, $-\text{SO}_2\text{NR}^1\text{R}^{1'}$, another aryl, another heteroaryl or another heterocyclyl and the like, whereby $\text{R}^{1'}$ has the meaning given above.

25

The term **heteroaryloxy** refers to a Het-O-group, wherein Het is a heteroaryl.

The term **sp³-hybridized** refers to a carbon atom and means that this carbon atom forms four bonds to four substituents placed in a tetragonal fashion around this carbon atom.

30

The expression **pharmaceutically acceptable** salts encompasses either salts with inorganic acids or organic acids like hydrochloric or hydrobromic acid, sulfuric acid, phosphoric acid, citric acid, formic acid, acetic acid, maleic acid, tartaric acid, benzoic acid, methanesulfonic acid, p-toluenesulfonic acid, and the like that
5 are non toxic to living organisms or in case the compound of formula I is acidic in nature with an inorganic base like an alkali or earth alkali base, e.g. sodium hydroxide, potassium hydroxide, calcium hydroxide and the like.

Compounds of the invention also include nitrosated compounds of the general
10 formula I that have been nitrosated through one or more sites such as oxygen (hydroxyl condensation), sulfur (sulffiydryl condensation) and/or nitrogen. The nitrosated compounds of the present invention can be prepared using conventional methods known to one skilled in the art. For example, known methods for nitrosating compounds are described in U.S. Pat. Nos. 5,380,758 and 5,703,073;
15 WO 97/27749; WO 98/19672; WO 98/21193; WO 99/00361 and Oae et al, Org. Prep. Proc. Int., 15(3):165-198 (1983), the disclosures of each of which are incorporated by reference herein in their entirety.

The compounds of the general formula I can contain two or more asymmetric
20 carbon atoms and may be prepared in form of optically pure enantiomers, mixtures of enantiomers such as racemates, diastereomers, mixtures of diastereomers, diastereomeric racemates, mixtures of diastereomeric racemates, and the meso-form and pharmaceutically acceptable salts thereof.

The present invention encompasses all these forms. Mixtures may be separated in
25 a manner known *per se*, i.e. by column chromatography, thin layer chromatography, HPLC or crystallization.

A group of preferred compounds are compounds of general formula I wherein Z,
Y, X, W, V, U, T, Q, L, and M are as defined in general formula I above and
30 wherein

k is 1

n is 0 and

m is 1.

Another group of preferred compounds of general formula I are those wherein Z,
5 Y, X, W, V, U, T, Q, M, k, m, and n are as defined in general formula I above and

L represents H; $-\text{COR}^{3''}$; $-\text{COOR}^{3''}$; $-\text{CONR}^{2''}\text{R}^{3''}$;

whereby $\text{R}^{2''}$ and $\text{R}^{3''}$ represent independently lower alkyl, lower cycloalkyl -
10 lower alkyl, which lower alkyl and lower cycloalkyl - lower alkyl groups are
unsubstituted or monosubstituted with halogen, cyano, hydroxy, $-\text{OCOCH}_3$,
 $-\text{CONH}_2$, $-\text{COOH}$ or $-\text{NH}_2$, with the proviso that a carbon atom is attached at the
most to one heteroatom in case this carbon atom is sp^3 -hybridized.

15 Another group of preferred compounds of general formula I above are those
wherein Z, Y, X, W, V, U, L, k, m, and n are as defined in general formula I and

T is $-\text{CONR}^1$;

Q is methylene;

20 M is aryl; heteroaryl.

Another group of also more preferred compounds of general formula I are those
wherein V, U, T, Q, M, L, k, m, and n are as defined in general formula I above
and

25

Z, Y, X and W represent $-\text{CH}-$.

Another group of also more preferred compounds of general formula I are those
wherein Z, Y, X, W, V, Q, T, M, L, k, m, and n are as defined in general formula I
30 above and

U is a mono-, di-, or trisubstituted phenyl or heteroaryl, wherein the substituents are halogen, lower alkyl, lower alkoxy, CF₃.

Especially preferred compounds of general formula I are those selected from the
5 group consisting of:

(*rac.*)-(1*R**, 5*S**)-(3-acetyl-7-{3-[2-(2-bromo-5-fluorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

10 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-chlorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-*tert*-butylphenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

15 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,3,6-trifluorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,5-difluorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

20

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-[3-(2-*o*-tolylloxyethyl)phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,3-dichlorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

25

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-chloro-5-methylphenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

30 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(3-chlorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

- (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[3-(2-bromo-5-fluorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;
- (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-chlorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;
- 5 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-*tert*-butylphenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;
- (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,3,6-trifluorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;
- 10 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,5-difluorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;
- (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-[3-(2-*o*-tolylloxypropyl)phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;
- 15 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,3-dichlorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;
- (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-chloro-5-methylphenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;
- 20 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(3-chlorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;
- 25 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(4-chlorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;
- (*rac.*)-(1*R**, 5*S**)-7-{3-[2-(2,6-dichloro-4-methylphenoxy)ethoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide;
- 30

(*rac.*)-(1*R**, 5*S**)-7-{3-[4-(2-fluoro-3-trifluoromethylphenoxy)butoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide;

5

(*rac.*)-(1*R**, 5*S**)-7-{3-[4-(2,6-dichloro-4-methylphenoxy)butoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide; and

10 (*rac.*)-(1*R**, 5*S**)-7-{3-[4-(2-chloro-6-fluoro-3-methylphenoxy)butoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide.

The compounds of general formula I and their pharmaceutically acceptable salts
15 may be used as therapeutics e.g. in form of pharmaceutical compositions. These pharmaceutical compositions containing at least one compound of general formula I and usual carrier materials and adjuvants may especially be used in the treatment or prophylaxis of disorders which are associated with a dysregulation of the renin angiotensin system (RAS), comprising cardiovascular and renal diseases.
20 Examples of such diseases are hypertension, congestive heart failure, pulmonary heart failure, coronary diseases, cardiac insufficiency, renal insufficiency, renal or myocardial ischemia, atherosclerosis, and renal failure. They can also be used to prevent restenosis after balloon or stent angioplasty, to treat erectile dysfunction, glomerulonephritis, renal colic, and glaucoma. Furthermore, they can be used in
25 the therapy and the prophylaxis of diabetic complications, complications after vascular or cardiac surgery, complications of treatment with immunosuppressive agents after organ transplantation, complications of cyclosporin treatment, as well as other diseases presently known to be related to the RAS.

30 In another embodiment, the invention relates to a method for the treatment and/or prophylaxis of diseases which are related to the RAS comprising hypertension, congestive heart failure, pulmonary hypertension, cardiac insufficiency, renal

insufficiency, renal or myocardial ischemia, atherosclerosis, renal failure, erectile dysfunction, glomerulonephritis, renal colic, glaucoma, diabetic complications, complications after vascular or cardiac surgery, restenosis, complications of treatment with immunosuppressive agents after organ transplantation, and other
5 diseases which are related to the RAS, which method comprises administering a compound according to general formula I to a human being or animal.

The invention further relates to the use of compounds of general formula I as defined above for the treatment and/or prophylaxis of disorders which are
10 associated with the Renin Angiotensin System (RAS) comprising hypertension, congestive heart failure, pulmonary hypertension, cardiac insufficiency, renal insufficiency, renal or myocardial ischemia, atherosclerosis, renal failure, erectile dysfunction, glomerulonephritis, renal colic, glaucoma, diabetic complications, complications after vascular or cardiac surgery, restenosis, complications of
15 treatment with immunosuppressive agents after organ transplantation, and other diseases known to be related to the RAS.

In addition, the invention relates to the use of compounds as defined above for the preparation of medicaments for the treatment and/or prophylaxis of diseases
20 which are associated with the RAS such as hypertension, coronary diseases, cardiac insufficiency, renal insufficiency, renal and myocardial ischemia, and renal failure.

The compounds of formula I may also be used in combination with one or more
25 other pharmacologically active compounds comprising ACE inhibitors, angiotensin II receptor antagonists, endothelin receptor antagonists, vasodilators, calcium antagonists, potassium activators, diuretics, sympatholitics, beta-adrenergic antagonists, alpha-adrenergic antagonists, and neutral endopeptidase inhibitors, for the treatment of the above-mentioned diseases or disorders.

30

All forms of prodrugs leading to an active component comprised by general formula I above are included in the present invention.


The compounds of general formula I can be manufactured by the methods outlined below, by the methods described in the examples or by analogous methods.

Preparation of the precursors:

Precursors are compounds which were prepared as key intermediates and/or building blocks and which were suitable for further transformations in parallel chemistry.


Bicyclononanone **A** was prepared from (4-benzyl-6-ethoxycarbonylmethyl-1-methylpiperazin-2-yl)acetic acid ethyl ester (Patent Application WO92/05174) as described in Scheme 1. Derivative **A** might also be present as enol form. In order to allow a coupling at the 7-position of bicyclononanone **A** with aryl bromides, the vinyl triflate derivative **B** was prepared.

Scheme 1



Compound **B** can be then transformed into compounds of type **C** by a *Negishi* coupling (Scheme 2), whereas R^b represents a side chain precursor suitable to construct the V-U-chain through one or several elementary chemical transformations. The R^b -substituent can be modified during the synthesis. After protecting group manipulations leading to compounds of type **D** the L^1 -substituent can be put in place (compounds of type **E**), whereas L^1 represents a substituent **L** as defined in general formula I, or a precursor of such a substituent. The ester functional group can be saponified to compounds of type **F**. After an amide coupling for instance precursors of type **G** can be obtained.

Scheme 2



- 5 Other type of compounds described in the general formula I can be prepared using the chemistry described in patent application WO03/093267.

Otherwise heterocyclic systems may be prepared according to the literature existing for similar compounds. For instance pyridine derivative H could be prepared from 2,6-dibromopyridine, wherein R^a is a substituent that may be easily transformed into a chain U-V as described in Formula I (Scheme 3; Bitman, R., et al.; *J. Org. Chem.*, 2000, 65, 7634). Pyridine J could be prepared by addition of a Grignard reagent on 4-bromopyridine, followed by oxidation (see Comins, D.; et al.; *J. Org. Chem.*, 1985, 50, 4410). From known 3,5-dibromopyridine a compound of type K could be prepared. Finally pyridines of type L could be prepared from commercially available 2-(pyridin-4-yl)-alcohol, according to the


10

15

literature described for similar compounds (Taylor, S. L.; et. al.; *J. Org. Chem.*, 1983, 48, 4156).

Scheme 3

5




Pyridine derivatives of type **M** could be prepared according to the literature (Scheme 4, Newkome, G. R.; et al.; *J. Am. Chem. Soc.*, 1975, 97, 3232). Known (4-bromopyridin-2-yl)oxy)acetic acid could lead to compound **N** by reduction. Compounds of type **O** could be prepared from 3,5-dibromopyridine, according to

10

Harrowven, D. C.; et al.; *Tetrahedron*, **2001**, 57, 4447. Known (2-bromopyridin-4-yl)acetic acid (Den Hertog, H. J.; et al.; *Chem. Pharm. Bull.*, **1975**, 23, 3008) could lead to compound P by reduction as well.

5

Scheme 4




A reduction of the known (4-bromopyrimidin-2-yl)acetic acid methyl ester (Brown, D. J.; et al.; *Australian J. Chem.*, **1978**, 31, 649) could lead to the desired pyrimidine intermediate Q (Scheme 5). As well, a palladium-catalyzed coupling could transform 2,4-dibromopyrimidine into a compound of type R (see

Gronowitz, S.; et al.; *Chem. Scripta*, **1986**, *26*, 305). Also, compounds of type S could be prepared from 6-alkyl-1H-pyrimidin-4-one.

Scheme 5


5



A palladium-catalyzed coupling of 2,4-dibromopyrimidine with bicyclononene **B** could lead to an intermediate of type **T**, which could be transformed into an intermediate of type **U** (Scheme 6). 6-Chloropyrazin-2-ylamine could be transformed in several steps into a pyrazine of type **V** (see Ghosh, A. K.; et al.; *J. Med. Chem.*, **1993**, *36*, 2300 or Jovanovic, M. V., *Heterocycles*, **1983**, *20*, 2011, or Hartman, G. D.; et al.; *J. Heterocyclic Chem.*; **1983**, *20*, 1089). Pyrimidines of type **W** could be prepared according to standard procedures. Finally a compounds

15 of type **X** can be prepared from 4,6-dichloropyrimidine.

Scheme 6





1) Protection of the amine


2) $\text{R}^a\text{CH}_2\text{OH}$

3) Deprotection of the amine

4) Diazotation and bromination


1) $\text{R}^a\text{CH}_2\text{OH}$ 2) NH_3

3) Diazotation, bromination



Pyridazinyl derivatives may be prepared as described in Scheme 7. For instance known 2-pyridazin-3-ylethanol **Y** (Rodriguez, L.; *et al.*; *Synlett*, 1990, 227) could be transformed into a pyridazinyl derivative **Z** following known chemistry (Sauer, J.; *et al.*; *Tetrahedron*, 1998, 54, 4297). Known 6-bromo-pyridazine-4-carboxylic acid ethyl ester **AA** (Dulayyi, A.; *et al.*; *Tetrahedron*, 1998, 54, 12897; Barlin, G. B.; *et al.*; *Australian J. Chem.*, 1977, 30, 2319) could be alkylated, then reduced to a derivative **AB** bearing the desired R^a-substituent. Also, known compound **AC** (Brundish, D. E.; *et al.*; *J. Labelled Compounds and Radiopharmaceuticals*, 1988, 25, 1371) could be transformed to derivative **AD** that could be reduced to pyridazinyl derivative **AE**. Hydrolysis could lead to derivative **AF**, finally O-alkylation to the desired intermediate **AG**.

Scheme 7




Preparation of final compounds

Precursors **G** were transformed into the corresponding aryl ethers (Scheme 8), using the *Mitsunobu* reaction conditions. After deprotection, the final compounds are obtained.

5

Scheme 8



- 10 The compounds of formula **I** and their pharmaceutically acceptable acid addition salts can be used as medicaments, e. g. in the form of pharmaceutical preparations for enteral, parenteral, or topical administration. They can be administered, for example, perorally, e. g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions, rectally, e. g. in the
- 15 form of suppositories, parenterally, e. g. in the form of injection solutions or infusion solutions, or topically, e. g. in the form of ointments, creams or oils.

The production of pharmaceutical preparations can be effected in a manner which will be familiar to any person skilled in the art by bringing the described

20 compounds of formula **I** and their pharmaceutically acceptable acid addition salts, optionally in combination with other therapeutically valuable substances, into a galenical administration form together with suitable, non-toxic, inert, therapeutically compatible solid or liquid carrier materials and, if desired, usual pharmaceutical adjuvants.

25

Suitable carrier materials are not only inorganic carrier materials, but also organic carrier materials. Thus, for example, lactose, corn starch or derivatives thereof, talc, stearic acid or its salts can be used as carrier materials for tablets, coated

tablets, dragées and hard gelatine capsules. Suitable carrier materials for soft gelatine capsules are, for example, vegetable oils, waxes, fats and semi-solid and liquid polyols (depending on the nature of the active ingredient no carriers are, however, required in the case of soft gelatine capsules). Suitable carrier materials
5 for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar and the like. Suitable carrier materials for injections are, for example, water, alcohols, polyols, glycerols and vegetable oils. Suitable carrier materials for suppositories are, for example, natural or hardened oils, waxes, fats and semi-liquid or liquid polyols. Suitable carrier materials for topical
10 preparations are glycerides, semi-synthetic and synthetic glycerides, hydrogenated oils, liquid waxes, liquid paraffins, liquid fatty alcohols, sterols, polyethylene glycols and cellulose derivatives.

Usual stabilizers, preservatives, wetting and emulsifying agents, consistency-
15 improving agents, flavour-improving agents, salts for varying the osmotic pressure, buffer substances, solubilizers, colorants and masking agents and antioxidants come into consideration as pharmaceutical adjuvants.

The dosage of compounds of formula I can vary within wide limits depending on
20 the disease to be controlled, the age and the individual condition of the patient and the mode of administration, and will, of course, be fitted to the individual requirements in each particular case. For adult patients a daily dosage of about 1 mg to about 1000 mg, especially about 50 mg to about 500 mg, comes into consideration.

25

The pharmaceutical preparations conveniently contain about 1 - 500 mg, preferably 5 - 200 mg of a compound of formula I.

The following examples serve to illustrate the present invention in more detail.
30 They are, however, not intended to limit its scope in any manner.

Examples

General remarks

- 5 The compounds were characterized at least by LC-MS and ^1H -NMR. Only the LC-MS data are given here.

Abbreviations

10	AcCl	Acetyl chloride
	ACE	Angiotensin Converting Enzyme
	AcOH	Acetic acid
	Ang	Angiotensin
	aq.	aqueous
15	Bn	Benzyl
	Boc	<i>tert</i> -Butyloxycarbonyl
	BSA	Bovine serum albumine
	BuLi	<i>n</i> -Butyllithium
	DIPEA	Diisopropylethylamine
20	DMAP	4- <i>N,N</i> -Dimethylaminopyridine
	DMSO	Dimethylsulfoxide
	EDC·HCl	Ethyl- <i>N,N</i> -dimethylaminopropylcarbodiimide hydrochloride
	EIA	Enzyme immunoassay
	eq.	equivalent
25	Et	Ethyl
	EtOAc	Ethyl acetate
	FC	Flash Chromatography
	HOBt	Hydroxybenzotriazol
	MeOH	Methanol
30	org.	organic
	PG	protecting group
	Ph	Phenyl

	RAS	Renin Angiotensin System
	RP18	Reversed phase column, filled with C ₁₈ hydrocarbon
	rt	room temperature
	sol.	Solution
5	TBAF	Tetrabutylammonium fluoride
	TBDMS	<i>tert</i> -Butyldimethylsilyl
	tBuOK	Potassium <i>tert</i> -butylate
	Tf	Trifluoromethylsulfonyl
	TFA	Trifluoroacetic acid
10	THF	Tetrahydrofuran
	TLC	Thin Layer Chromatography
	TMAD	<i>N,N,N',N'</i> -Tetramethylazodicarboxamide

Preparation of the precursors

15

[4-(3-Bromophenoxy)butoxy]-*tert*-butyldimethylsilane

A mixture of 3-bromophenol (10.0 g, 56.6 mmol), KI (50 mg) and K₂CO₃ (12.5 g, 90.6 mmol) in DMF (100 mL) was stirred at 85 °C for 30 min. 4-Bromobutanol
20 (8.75 g, 57.2 mmol) was added and the mixture was stirred at 85 °C for 16 h. 4-Bromobutanol (8.75 g, 57.2 mmol) was added again and the mixture was stirred at 85 °C overnight. The mixture was poured onto a mixture of ice and water (500 mL), and extracted with CH₂Cl₂. The combined org. extracts were washed with aq. 1M NaOH, dried over MgSO₄, filtered, and the solvents were removed under
25 reduced pressure. Purification by FC yielded 4-(3-bromophenoxy)-butan-1-ol (4.66 g, 34%).

This compound was dissolved in DMF (75 mL), and imidazol (3.27 g, 47.5 mmol), then TBDMS-Cl (3.02 g, 19.0 mmol) were added. The mixture was stirred at rt overnight. The mixture was poured onto a sol. of NaHCO₃ (0.5 g) in
30 water (125 mL). The resulting mixture was extracted with Et₂O (2x). The combined org. extracts were dried over MgSO₄, filtered, and the solvents were

removed under reduced pressure. Purification by FC yielded the title compound (6.78 g, quantitative yield).

(rac.)-(1*R, 5*S**)-9-Methyl-7-oxo-3,9-diazabicyclo[3.3.1]nonane-3,6-dicarboxylic acid 3-*tert*-butyl ester 6-ethyl ester (A)**

(4-Benzyl-6-ethoxycarbonylmethyl-1-methyl-piperazin-2-yl)acetic acid ethyl ester (Patent WO 92/05174) (71.0 g, 0.196 mol) was dissolved in MeOH (400 mL). TFA (77.8 mL, 1.02 mol) was added and the flask was purged with nitrogen.

10. Pd/C (10%, 50% moisture, 3.6 g) was added. The flask was closed and purged with hydrogen (3x). After 1 day, the mixture was filtered through *Celite* and washed with MeOH. The solvents were removed under reduced pressure and the foamy residue (92.7 g) was dried under high vacuum. A sol. of *t*BuOK (117.2 g, 1.04 mol) in toluene (3.07 L) was heated to reflux. A sol. of the crude piperazine

15. formerly obtained, dissolved in THF (300 mL), was added dropwise over 50 min. The black mixture was stirred for 10 further min. and allowed to cool to rt. The mixture was cooled to 0 °C and AcOH (36.6 mL, 0.635 mol) was added. The solvents were removed under reduced pressure. This crude material was suspended in CH₂Cl₂ (400 mL) and cooled to 0 °C. DIPEA (19.1 mL, 112 mmol)

20. was added. A sol. of Boc₂O (24.3 g, 113 mmol) in CH₂Cl₂ (200 mL) was added dropwise. The mixture was stirred for 1 h at 0 °C, then 1 h at rt. The mixture was washed with aq. 10% Na₂CO₃ (2x). The org. extracts were dried over MgSO₄, filtered and the solvents were evaporated under reduced pressure. The residue was purified by FC (EtOAc/heptane 1:1 → EtOAc). The title compound was

25. obtained as oil (24.5 g, 38%). *R*_f = 0.05 (EtOAc/heptane 1:1) or 0.56 (MeOH/CH₂Cl₂ 1:9). LC-MS: *R*_t = 2.94; ES⁺: 325.19.

(rac.)-(1*R, 5*S**)-9-Methyl-7-trifluoromethanesulfonyloxy-3,9-diazabicyclo[3.3.1]non-6-ene-3,6-dicarboxylic acid 3-*tert*-butyl ester 6-ethyl ester (B)**

30

A sol. of bicyclononanone A (2.22 g, 6.80 mmol) in THF (50 mL) was cooled to 0 °C and NaH (about 60% in mineral oil, 326 mg, about 8.2 mmol) was added. A

gas evolution was observed. After 20 min, Ti_2NPh (3.22 g, 9.00 mmol) was added. 10 min later, the ice bath was removed. After 3 h, the sol. was diluted with EtOAc and washed with brine (1x). The org. extracts were dried over MgSO_4 , filtered, and the solvents were removed under reduced pressure.

5 Purification by FC (EtOAc/heptane 3:1 \rightarrow EtOAc) yielded the title compound as an oil (2.50 g, 80%). $R_f = 0.15$ (EtOAc/heptane 1:1). LC-MS: $R_t = 4.73$; ES+: 458.95.

(rac.)-(1R*, 5S*)-7-{3-[2-(*tert*-Butyldimethylsilanyloxy)ethyl]phenyl}-9-methyl-3,9-diazabicyclo[3.3.1]non-6-ene-3,6-dicarboxylic acid 3-*tert*-butyl ester 6-ethyl ester (C1)

10

BuLi (1.6M in hexane, 8.52 mL, 13.6 mmol) was added to a sol. of [2-(3-bromophenyl)ethoxy]-*tert*-butyldimethylsilane (4.28 g, 13.6 mmol, prepared by

15 silylation of 2-(3-bromophenyl)ethanol) in THF (60 mL) at -78°C . The mixture was stirred for 30 min at this temperature, and ZnCl_2 (1M in THF, prepared from ZnCl_2 dried at 160°C for 3 h and THF, 15.9 mL, 15.9 mmol) was added. The mixture was allowed to warm to rt and a sol. of bicyclononene **B** (2.50 g, 5.45 mmol) in THF (15 mL), then $\text{Pd}(\text{PPh}_3)_4$ (147 mg, 0.13 mmol) were added. The

20 mixture was heated to 40°C and stirred at this temperature for 30 min. The mixture was allowed to cool to rt and aq. 1M HCl (2 mL) was added. The mixture was diluted with EtOAc and washed with aq. 1M NaOH (1x). The org. extracts were dried over MgSO_4 , filtered, and the solvents were removed under reduced pressure. Purification by FC (MeOH/ CH_2Cl_2 10:490 \rightarrow 15:485 \rightarrow 20:480 \rightarrow

25 25:475 \rightarrow 50:450) yielded the title compound (2.41 g, 80%). LC-MS; $R_t = 5.02$ ES+: 545.31.

(rac.)-(1R*, 5S*)-7-{3-[2-(*tert*-Butyldimethylsilanyloxy)propyl]phenyl}-9-methyl-3,9-diazabicyclo[3.3.1]non-6-ene-3,6-dicarboxylic acid 3-*tert*-butyl ester 6-ethyl ester (C2)

30

BuLi (1.6M in hexane, 10.22 mL, 16.4 mmol) was added to a sol. of [2-(3-bromophenyl)propoxy]-*tert*-butyldimethylsilane (4.95 g, 16.3 mmol, prepared by silylation of 2-(3-bromophenyl)propanol, P. Beak, *et al.*, *J. Org. Chem.*, **1989**, *54*, 5574) in THF (60 mL) at -78 °C. The mixture was stirred for 30 min at this
5 temperature, and ZnCl₂ (1M in THF, prepared from ZnCl₂ dried at 160 °C for 3 h and THF, 19.0 mL, 19.0 mmol) was added. The mixture was allowed to warm to rt and a sol. of bicyclononene B (3.00 g, 6.54 mmol) in THF (15 ml), then Pd(PPh₃)₄ (189 mg, 0.16 mmol) were added. The mixture was heated to 40 °C and stirred at this temperature for 30 min. The mixture was allowed to cool to rt
10 and aq. 1M HCl (2 mL) was added. The mixture was diluted with EtOAc and washed with aq. 1M NaOH (1x). The org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification by FC (MeOH/CH₂Cl₂ 10:490 → 15:485 → 20:480 → 25:475 → 50:450) yielded the title compound (2.74 g, 75%). LC-MS; R_t = 5.10 ES+: 559.32.

15

(rac)-(1R*, 5S*)-7-{3-[2-(*tert*-Butyldimethylsilyloxy)ethoxy]phenyl}-9-methyl-3,9-diazabicyclo[3.3.1]non-6-ene-3,6-dicarboxylic acid 3-*tert*-butyl ester 6-ethyl ester (C3)

20 BuLi (1.6M in hexane, 19.2 mL, 30.7 mmol) was added to a sol. of [2-(4-bromophenoxy)ethoxy]-*tert*-butyldimethylsilane (9.40 g, 28.4 mmol, Morita, C.; *et al.*, *Heterocycles*, **2000**, *52*, 1163) in THF (55 mL) at -78 °C. The mixture was stirred for 30 min at this temperature, and ZnCl₂ (1M in THF, prepared from ZnCl₂ dried at 160 °C for 3 h and THF, 35.5 mL, 35.5 mmol) was added. The mixture was
25 allowed to warm to rt and a sol. of bicyclononene B (10.8 g, 23.6 mmol) in THF (20 ml), then Pd(PPh₃)₄ (0.70 g, 0.59 mmol) were added. The mixture was heated to 40 °C and stirred at this temperature for 30 min. The mixture was allowed to cool to rt and aq. 1M HCl (2 mL) was added. The mixture was diluted with EtOAc and washed with aq. 1M NaOH (1x). The org. extracts were dried over
30 MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification by FC (MeOH/CH₂Cl₂ 10:490 → 15:485 → 20:480 → 25:475 →

50:450) yielded the title compound (11.7 g, 89%). LC-MS; R_t = 0.98 ES+: 561.10.

(rac.)-(1*R, 5*S**)-7-{3-[4-(*tert*-Butyldimethylsilyloxy)butoxy]phenyl}-9-methyl-3,9-diazabicyclo[3.3.1]non-6-ene-3,6-dicarboxylic acid 3-*tert*-butyl ester 6-ethyl ester (C4)**

BuLi (1.6M in hexane, 12.75 mL, 20.4 mmol) was added to a sol. of [4-(3-bromophenoxy)butoxy]-*tert*-butyldimethylsilane (6.78 g, 18.8 mmol) in THF (100 mL) at -78 °C. The mixture was stirred for 30 min at this temperature, and ZnCl₂ (1M in THF, prepared from ZnCl₂ dried at 160 °C for 3 h and THF, 23.6 mL, 23.6 mmol) was added. The mixture was allowed to warm to rt and a sol. of bicyclononene **B** (7.2 g, 15.7 mmol) in THF (20 ml), then Pd(PPh₃)₄ (0.468 g, 0.393 mmol) were added. The mixture was stirred at rt for 1 h. The mixture was allowed to cool to rt and aq. 1M HCl (2 mL) was added. The mixture was diluted with EtOAc and washed with aq. 1M NaOH (1x). The org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification by FC (MeOH/CH₂Cl₂ 10:490 → 15:485 → 20:480 → 25:475 → 50:450) yielded the title compound (2.60 g, 28%). LC-MS; R_t = 1.02 ES+: 589.10.

(rac.)-(1*R, 5*S**)-7-{3-[2-(*tert*-Butyldimethylsilyloxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-3,6,9-tricarboxylic acid 3-*tert*-butyl ester 6-ethyl ester 9-(2,2,2-trichloro-1,1-dimethylethyl) ester (D1)**

25

A sol. of bicyclononene **C1** (2.3 g, 4.22 mmol) and β,β,β-trichloro-*tert*-butylchloroformate (10.4 g, 43.3 mmol) in CH₂ClCH₂Cl (40 ml) was heated to reflux for 3 h. The mixture was allowed to cool to rt and the solvents were removed under reduced pressure. Purification of the residue by FC (EtOAc / heptane 1:9 → 1:4 → 2:3) yielded the title compound (2.73 g, 88%). LC-MS: R_t = 8.00 min.

30

(rac.)-(1*R, 5*S**)-7-[3-[2-(*tert*-Butyldimethylsilanyloxy)propyl]phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-3,6,9-tricarboxylic acid 3-*tert*-butyl ester 6-ethyl ester 9-(2,2,2-trichloro-1,1-dimethylethyl) ester (D2)**

- 5 A sol. of bicyclononene C2 (2.74 g, 4.90 mmol) and β,β,β -trichloro-*tert*-butylchloroformate (11.76 g, 49.0 mmol) in $\text{CH}_2\text{ClCH}_2\text{Cl}$ (50 ml) was heated to reflux for 3 h. The mixture was allowed to cool to rt and the solvents were removed under reduced pressure. Purification of the residue by FC (EtOAc / heptane 1:9 \rightarrow 1:4 \rightarrow 2:3) yielded the title compound (3.01 g, 85%). LC-MS: R_t
10 = 8.05 min.

(rac.)-(1*R, 5*S**)-7-[3-(2-Hydroxyethoxy)phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-3,6,9-tricarboxylic acid 3,9-di-*tert*-butyl ester 6-ethyl ester (D3)**

- 15 NaHCO_3 (17.4 g, 208 mmol) and 1-chloroethyl chloroformate (22.4 mL, 208 mmol) were added to a sol. of bicyclononene C3 (11.65 g, 20.8 mmol) in 1,2-dichloroethane (180 mL). The sol. was heated to reflux. After 4.5 h, the reaction mixture was allowed to cool to rt, and the solvents were removed under reduced pressure. MeOH (180 mL) was added. The mixture was stirred at rt for 30 min,
20 and the solvents were removed under reduced pressure. The residue was dissolved in CH_2Cl_2 (220 mL), DIPEA (17.8 mL, 104 mmol) was added, and the mixture was cooled to 0 °C. Boc_2O (13.6 g, 62.4 mmol) was added and the mixture was stirred at rt for 2 h. The mixture was washed with aq. 1M HCl (1x), and aq. sat. NaHCO_3 (1x). The org. extracts were dried over MgSO_4 , filtered, and the
25 solvents were removed under reduced pressure. Purification of the residue by FC yielded the title compound (9.57 g, 87%). LC-MS: R_t = 1.02 min; ES+: 477.02.

(rac.)-(1*R, 5*S**)-7-[3-(4-Hydroxybutoxy)phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-3,6,9-tricarboxylic acid 3,9-di-*tert*-butyl ester 6-ethyl ester (D4)**

30

NaHCO_3 (3.74 g, 44.2 mmol) and 1-chloroethyl chloroformate (4.85 mL, 44.2 mmol) were added to a sol. of bicyclononene C4 (2.60 g, 4.41 mmol) in 1,2-

dichloroethane (30 mL). The sol. was heated to reflux. After 3 h, the reaction mixture was allowed to cool to rt, and the solvents were removed under reduced pressure. MeOH (30 mL) was added. The mixture was stirred at rt for 30 min, and the solvents were removed under reduced pressure. The residue was dissolved in CH₂Cl₂ (50 mL), DIPEA (3.75 mL, 22.1 mmol) was added, and the mixture was cooled to 0 °C. Boc₂O (2.95 g, 13.2 mmol) was added and the mixture was stirred at rt for 2 h. The mixture was washed with aq. 1M HCl (1x), and aq. sat. NaHCO₃ (1x). The org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification of the residue by FC yielded the title compound (1.71 g, 69%). LC-MS: R_t = 1.05 min; ES+: 505.08.

(rac.)-(1*R, 5*S**)-3-Acetyl-7-[3-(2-hydroxyethyl)phenyl]-3,9-diazabicyclo-[3.3.1]non-6-ene-6,9-dicarboxylic acid 6-ethyl ester 9-(2,2,2-trichloro-1,1-dimethylethyl) ester (E1)**

15

HCl (4M in dioxane, 30 mL) was added to a sol. of bicyclononene **D1** (2.73 g, 3.72 mmol) in CH₂Cl₂ (30 mL) at 0 °C. The mixture was stirred at 0 °C for 20 min, then at rt for 2.5 h. The solvents were evaporated under reduced pressure and the residue was dried under high vacuum. A mixture of this residue, DMAP (23 mg, 0.19 mmol,) and DIPEA (2.54 mL, 15.3 mmol) in THF (15 mL) was cooled to -78 °C. AcCl (0.29 mL, 3.70 mmol) was added slowly. The mixture was stirred for 30 min, and MeOH (2 mL) was added. The mixture was allowed to warm up to rt and was diluted with EtOAc, washed with aq. 1M HCl (2x) and aq. sat. NaHCO₃ (1x). The org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification of the residue by FC (EtOAc / heptane 1:1 → EtOAc → EtOAc / MeOH 1:9) yielded the title compound (1.25 g, 60%). LC-MS: R_t = 6.62.

(rac.)-(1*R, 5*S**)-3-Acetyl-7-[3-(2-hydroxypropyl)phenyl]-3,9-diazabicyclo-[3.3.1]non-6-ene-6,9-dicarboxylic acid 6-ethyl ester 9-(2,2,2-trichloro-1,1-dimethylethyl) ester (E2)**

30

HCl (4M in dioxane, 30 mL) was added to a sol. of bicyclononene **D2** (3.01 g, 4.02 mmol) in CH₂Cl₂ (30 ml) at 0 °C. The mixture was stirred at 0 °C for 20 min, then at rt for 2.5 h. The solvents were evaporated under reduced pressure and the residue was dried under high vacuum. A mixture of this residue, DMAP (25 mg, 0.20 mmol,) and DIPEA (2.75 ml, 16 mmol) in THF (20 mL) was cooled to -78 °C. AcCl (0.31 mL, 4.10 mmol) was added slowly. The mixture was stirred for 30 min, and MeOH (2 mL) was added. The mixture was allowed to warm up to rt and was diluted with EtOAc, washed with aq. 1M HCl (2x) and aq. sat. NaHCO₃ (1x). The org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification of the residue by FC (EtOAc / heptane 1:1 → EtOAc → EtOAc / MeOH 1:9) yielded the title compound (1.56 g, 69%).

(*rac.*)-(1*R**, 5*S**)-3-Acetyl-7-[3-(2-hydroxyethyl)phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-6,9-dicarboxylic acid 9-(2,2,2-trichloro-1,1-dimethylethyl) ester (**F1**)

A mixture of bicyclononene **E1** (1.25 g, 2.22 mmol) in EtOH (56 ml) and aq. 1M NaOH (56 mL) was stirred at 80 °C for 2.5 h. The mixture was allowed to cool to rt and acidified with aq 1M HCl. The mixture was extracted with EtOAc (3x) and the org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification of the crude by FC (EtOAc → MeOH/EtOAc 1:19 → 1:9) yielded the title compound (360 mg 35%). LC-MS: R_t = 4.66; MS-: 530.82.

25

(*rac.*)-(1*R**, 5*S**)-3-Acetyl-7-[3-(2-hydroxypropyl)phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-6,9-dicarboxylic acid 9-(2,2,2-trichloro-1,1-dimethylethyl) ester (**F2**)

A mixture of bicyclononene **E2** (1.56 g, 2.70 mmol) in EtOH (70 ml) and aq. 1M NaOH (70 mL) was stirred at 80 °C for 2.5 h. The mixture was allowed to cool to rt and acidified with aq 1M HCl. The mixture was extracted with EtOAc (3x) and

the org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification of the crude by FC (EtOAc → MeOH/EtOAc 1:19 → 1:9) yielded the title compound (1.18 g 80%).

5 **(rac.)-(1*R**, 5*S**)-7-[3-(*tert*-Butyldimethylsilanyloxy)ethoxyphenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-3,6,9-tricarboxylic acid 3,9-di-*tert*-butyl ester (F3)**

A mixture of bicyclononene **D3** (6.27 g, 11.8 mmol) in EtOH (210 ml) and aq. 1M NaOH (90 mL) was stirred at 80 °C for 5.5 h. The mixture was allowed to cool to
10 rt and acidified with aq 1M HCl. The mixture was extracted with EtOAc (3x) and the org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. The crude compound (5.21 g 10.4 mmol), imidazol (2.12 g, 31.2 mmol) and TBDMS-Cl (3.45 g, 22.9 mmol) in DMF (50 mL) were stirred
15 at rt overnight. Aq. sat. NH₄Cl was added and the mixture was extracted with hexane (3x). The combined org. extracts were dried over MgSO₄, filtered, and the solvents were evaporated under reduced pressure. A mixture of this crude product and K₂CO₃ (0.6 g) in THF (30 ml), MeOH (10 ml), and H₂O (10 ml) was stirred at
20 rt for 3 h. Aq. sat. NH₄Cl was added and this mixture was extracted with Et₂O (3x). The combined org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. This yielded the title compound (6.21 g, 85%) that was used without further purification.

25 **(rac.)-(1*R**, 5*S**)-7-{3-[4-(*tert*-Butyldimethylsilanyloxy)butoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-3,6,9-tricarboxylic acid 3,9-di-*tert*-butyl ester (F4)**

A mixture of bicyclononene **D4** (1.71 g, 3.05 mmol) in EtOH (35 ml) and aq. 1M NaOH (15 mL) was stirred at 80 °C for 4 h. The mixture was allowed to cool to rt
30 and acidified with aq 1M HCl. The mixture was extracted with EtOAc (3x) and the org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. The crude compound (1.65 g 3.05 mmol), imidazol (0.85

g, 12.4 mmol) and TBDMS-Cl (2.50 g, 7.7 mmol) in DMF (35 mL) were stirred at rt overnight. Aq. sat. NH₄Cl was added and the mixture was extracted with hexane (3x). The combined org. extracts were dried over MgSO₄, filtered, and the solvents were evaporated under reduced pressure. A mixture of this crude product and K₂CO₃ (0.6 g) in THF (30 ml), MeOH (10 ml), and H₂O (10 ml) was stirred at rt for 3 h. Aq. sat. NH₄Cl was added and this mixture was extracted with Et₂O (3x). The combined org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. This yielded the title compound (1.79 g, 87%) that was used without further purification. LC-MS: R_t = 1.35; ES+: 783.31.

(rac.)-(1R*, 5S*)-3-Acetyl-7-[3-(2-hydroxyethyl)phenyl]-6-(methyl-phenethyl-carbamoyl)-3,9-diazabicyclo[3.3.1]non-6-ene-9-carboxylic acid 2,2,2-trichloro-1,1-dimethylethyl ester (G1)

A mixture of bicyclononene F1 (360 mg, 0.67 mmol), methylphenethylamine (0.28 mL, 2.01 mmol), DMAP (20 mg, 0.17 mmol), DIPEA (0.74 mL, 4.30 mmol), HOBt (22 mg, 0.16 mmol) and EDC·HCl (310 mg, 1.68 mmol) in CHCl₃ (10 mL) was stirred for 3 days. The mixture was diluted in CH₂Cl₂ and washed with aq. 1M HCl (2x), and aq. sat. NaHCO₃ (1x). The org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification by FC (EtOAc / heptane 1:4 → 1:1 → 4:1 → EtOAc → MeOH/EtOAc 1:19 → 90:10) yielded the title compound (350 mg, 80%). LC-MS: R_t = 5.37; ES+: 650.19.

(rac.)-(1R*, 5S*)-3-Acetyl-7-[3-(2-hydroxypropyl)phenyl]-6-(methyl-phenethylcarbamoyl)-3,9-diazabicyclo[3.3.1]non-6-ene-9-carboxylic acid 2,2,2-trichloro-1,1-dimethylethyl ester (G2)

A mixture of bicyclononene F2 (790 mg, 1.44 mmol), methylphenethylamine (0.63 mL, 4.33 mmol), DMAP (44 mg, 0.36 mmol), DIPEA (0.99 mL, 5.94 mmol), HOBt (48 mg, 0.36 mmol) and EDC·HCl (690 mg, 3.60 mmol) in CHCl₃

(20 mL) was stirred for 3 days. The mixture was diluted in CH₂Cl₂ and washed with aq. 1M HCl (2x), and aq. sat. NaHCO₃ (1x). The org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification by FC (EtOAc / heptane 1:4 → 1:1 → 4:1 → EtOAc →
5 MeOH/EtOAc 1:19 → 90:10) yielded the title compound (750 mg, 78%). LC-MS: R_t = 5.43; ES+: 664.13.

(rac.)-(1R*, 5S*)-6-[Cyclopropyl-(3-methoxy-2-methylbenzyl)carbamoyl]-7-[3-(2-hydroxyethoxy)phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-3,9-dicarboxylic acid di-*tert*-butyl ester (G3)
10

(1-Chloro-2-methylpropenyl)dimethylamine (1.50 mL, 11.6 mmol) was added to a sol. of compound F3 (6.5 g, 10.5 mmol) in CH₂Cl₂ (100 mL) at rt. The mixture was stirred for 20 min. Cyclopropyl-(3-methoxy-2-methylbenzyl)amine (prepared
15 by reductive amination from 3-methoxy-2-methylbenzaldehyde, Comins, D. L.; Brown, J. D., *J. Org. Chem.*, **1989**, *54*, 3730 and cyclopropylamine, 2.01 g, 10.5 mmol), and DIPEA (11.0 mL, 31.5 mmol) were added. The mixture was stirred for 35 min, and aq. 10% citric acid was added. The phases were separated, and the org. phase was washed with aq. 10% citric acid. The combined aq. phases
20 were extracted back with CH₂Cl₂. The combined org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. The crude title compound (8.30 g, 10.5 mmol) was dissolved in THF (170 mL) and TBAF (3.70 g, 11.5 mmol) was added. The mixture was stirred for 25 min. The solvents were removed under reduced pressure, and the residue was diluted with
25 EtOAc. The mixture was washed with water, and brine. The org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification of the residue by FC yielded the title compound (4.69 g, 66%). LC-MS: R_t = 1.07; ES+: 678.32.

(*rac.*)-(1*R**, 5*S**)-6-[Cyclopropyl-(3-methoxy-2-methylbenzyl)carbamoyl]-7-[3-(4-hydroxybutoxy)phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-3,9-dicarboxylic acid di-*tert*-butyl ester (G4)

- 5 (1-Chloro-2-methylpropenyl)dimethylamine (0.42 mL, 3.04 mmol) was added to a sol. of compound F4 (1.79 g, 2.76 mmol) in CH₂Cl₂ (50 mL) at rt. The mixture was stirred for 30 min. Cyclopropyl-(3-methoxy-2-methylbenzyl)amine (prepared by reductive amination from 3-methoxy-2-methylbenzaldehyde, Comins, D. L.; Brown, J. D., *J. Org. Chem.*, **1989**, 54, 3730 and cyclopropylamine, 0.53 g, 2.76
- 10 mmol), and DIPEA (1.44 mL, 8.30 mmol) were added. The mixture was stirred for 1 h, and aq. 10% citric acid was added. The phases were separated, and the org. phase was washed with aq. 10% citric acid. The combined aq. phases were extracted back with CH₂Cl₂. The combined org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. The crude title
- 15 compound (2.29 g) was dissolved in THF (50 mL) and TBAF (1.82 g, 5.58 mmol) was added. The mixture was stirred for 2 h. The solvents were removed under reduced pressure, and the residue was diluted with EtOAc. The mixture was washed with water, and brine. The org. extracts were dried over MgSO₄, filtered, and the solvents were removed under reduced pressure. Purification of the
- 20 residue by FC yielded the title compound (1.22 g, 62%). LC-MS: R_t = 1.09; ES+: 706.28.

Preparation of the final compounds

25 *General procedure (A) for the formation of aryl ether (Mitsunobu reaction)*

- The precursor (0.05 mmol) was dissolved or suspended in toluene (1.00 mL). The phenol derivative (0.075 mmol) in toluene (0.50 mL) was added. TMAD (0.075 mmol) in toluene (0.50 mL) was added, followed by tributylphosphine (0.15
- 30 mmol). The reaction mixture was stirred for 2 h at rt and then 2 h at 60 °C. Sometimes, it was necessary to add a second portion of tributylphosphine and to stir overnight. Sometimes, THF was necessary as cosolvent to dissolve the

reactants. The reaction mixture was allowed to cool to rt, then water was added. The mixture was extracted with EtOAc, and the org. extracts were evaporated under reduced pressure.

5 ***General procedure (B) for the cleavage of the 2,2,2-trichloro-1,1-dimethylethylcarbamate protecting group:***

The crude product from another general procedure was dissolved in THF/AcOH (1:0.1) and treated with zinc (20 eq.). The suspension was stirred for 5 h and
10 filtered through celite, which was washed with EtOAc. The filtrate was evaporated under reduced pressure and the residue was purified by HPLC.

Examples:

15 **Example 1**

(rac.)-(1R*, 5S*)-3-Acetyl-7-{3-[2-(2-chlorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt

According to the general procedures A and B, starting from compound G1 and 2-
20 chlorophenol. LC-MS: R_t = 0.83 min, ES⁺ = 558.11.

Example 2

***(rac.)-(1R*, 5S*)-3-Acetyl-7-{3-[2-(2-tert-butylphenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide
25 formate salt***

According to the general procedures A and B, starting from compound G1 and 2-
tert-butylphenol. LC-MS: R_t = 0.93 min, ES⁺ = 580.20.

Example 3

(rac.)-(1*R, 5*S**)-3-Acetyl-7-{3-[2-(2,3,6-trifluorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

5

According to the general procedures A and B, starting from compound G1 and 2,3,6-trifluorophenol. LC-MS: $R_t = 0.82$ min, $ES^+ = 578.11$.

Example 4

10 **(rac.)-(1*R**, 5*S**)-3-Acetyl-7-{3-[2-(2,5-difluorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

15 According to the general procedures A and B, starting from compound G1 and 2,5-difluorophenol. LC-MS: $R_t = 0.81$ min, $ES^+ = 560.11$.

Example 5

20 **(rac.)-(1*R**, 5*S**)-3-Acetyl-7-[3-(2-*o*-tolylloxyethyl)phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

According to the general procedures A and B, starting from compound G1 and 2-methylphenol. LC-MS: $R_t = 0.84$ min, $ES^+ = 538.15$.

Example 6

25 **(rac.)-(1*R**, 5*S**)-3-Acetyl-7-{3-[2-(2,3-dichlorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

30 According to the general procedures A and B, starting from compound G1 and 2,3-dichlorophenol. LC-MS: $R_t = 0.87$ min, $ES^+ = 592.06$.

Example 7

(rac.)-(1*R, 5*S**)-3-Acetyl-7-{3-[2-(2-chloro-5-methylphenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

5

According to the general procedures A and B, starting from compound G1 and 2-chloro-5-methylphenol. LC-MS: $R_t = 0.87$ min, $ES^+ = 572.11$.

Example 8

10 **(rac.)-(1*R**, 5*S**)-3-Acetyl-7-{3-[2-(3-chlorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

According to the general procedures A and B, starting from compound G1 and 3-chlorophenol. LC-MS: $R_t = 0.85$ min, $ES^+ = 558.11$.

15

Example 9

(rac.)-(1*R, 5*S**)-(3-Acetyl-7-{3-[2-(2-bromo-5-fluorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

20

According to the general procedures A and B, starting from compound G1 and 2-bromo-5-fluorophenol.

Example 10

25 **(rac.)-(1*R**, 5*S**)-3-Acetyl-7-{3-[3-(2-bromo-5-fluorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

30 According to the general procedures A and B, starting from compound G2 and 2-bromo-5-fluorophenol. LC-MS: $R_t = 0.89$ min, $ES^+ = 636.03$.

Example 11

(rac.)-(1*R, 5*S**)-3-Acetyl-7-{3-[2-(2-chlorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

5

According to the general procedures A and B, starting from compound G2 and 2-chlorophenol. LC-MS: $R_t = 0.86$ min, $ES^+ = 572.10$.

Example 12

10 **(rac.)-(1*R**, 5*S**)-3-Acetyl-7-{3-[2-(2-*tert*-butylphenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

According to the general procedures A and B, starting from compound G2 and 2-*tert*-butylphenol. LC-MS: $R_t = 0.97$ min, $ES^+ = 594.23$.

15**Example 13**

(rac.)-(1*R, 5*S**)-3-Acetyl-7-{3-[2-(2,3,6-trifluorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

20

According to the general procedures A and B, starting from compound G2 and 2,3,6-trifluorophenol. LC-MS: $R_t = 0.85$ min, $ES^+ = 592.13$.

25 **Example 14**

(rac.)-(1*R, 5*S**)-3-Acetyl-7-{3-[2-(2,5-difluorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

30 According to the general procedures A and B, starting from compound G2 and 2,5-difluorophenol. LC-MS: $R_t = 0.84$ min, $ES^+ = 574.14$.

Example 15

(rac.)-(1*R, 5*S**)-3-Acetyl-7-[3-(2-*o*-tolylloxypropyl)phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

5 According to the general procedures A and B, starting from compound **G2** (50 mg) and 2-methylphenol. LC-MS: $R_t = 0.88$ min, $ES^+ = 552.17$.

Example 16

10 **(rac.)-(1*R**, 5*S**)-3-Acetyl-7-[3-[2-(2,3-dichlorophenoxy)propyl]phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

According to the general procedures A and B, starting from compound **G2** and 2,3-dichlorophenol. LC-MS: $R_t = 0.91$ min, $ES^+ = 606.08$.

15

Example 17

(rac.)-(1*R, 5*S**)-3-Acetyl-7-[3-[2-(2-chloro-5-methylphenoxy)propyl]phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

20

According to the general procedures A and B, starting from compound **G2** and 2-chloro-5-methylphenol. LC-MS: $R_t = 0.90$ min, $ES^+ = 586.12$.

Example 18

25 **(rac.)-(1*R**, 5*S**)-3-Acetyl-7-[3-[2-(3-chlorophenoxy)propyl]phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt**

30 According to the general procedures A and B, starting from compound **X2** (50 mg) and 3-chlorophenol. LC-MS: $R_t = 0.88$ min, $ES^+ = 572.12$.

Example 19

(*rac.*)-(1*R**, 5*S**)-3-Acetyl-7-{3-[2-(4-chlorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide formate salt

5

According to the general procedures A and B, starting from compound G2 and 4-chlorophenol. LC-MS: R_t = 0.88 min, ES+ = 572.11.

Example 20

10 (*rac.*)-(1*R**, 5*S**)-7-{3-[2-(2,6-Dichloro-4-methylphenoxy)ethoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide

15 According to the general procedures A and B, starting from compound G3 and 2,6-dichloro-4-methylphenol. LC-MS: R_t = 0.88 min, ES+ = 572.11.

Example 21

20 (*rac.*)-(1*R**, 5*S**)-7-{3-[4-(2-Fluoro-3-trifluoromethylphenoxy)butoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide

According to the general procedures A and B, starting from compound G4 and 2-fluoro-3-trifluoromethylphenol. LC-MS: R_t = 0.88 min, ES+ = 572.11.

Example 22

25 (*rac.*)-(1*R**, 5*S**)-7-{3-[4-(2,6-Dichloro-4-methylphenoxy)butoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide

30 According to the general procedures A and B, starting from compound G4 and 2,6-dichloro-4-methylmethoxyphenol. LC-MS: R_t = 0.88 min, ES+ = 572.11.

Example 23

(*rac.*)-(1*R**, 5*S**)-7-{3-[4-(2-Chloro-6-fluoro-3-methylphenoxy)butoxy]-phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide

5

According to the general procedures A and B, starting from compound G4 and 2-chloro-6-fluoro-3-methylphenol. LC-MS: $R_t = 0.88$ min, $ES^+ = 572.11$.

10 The following assay was carried out in order to determine the activity of the compounds of general formula I and their salts.

Inhibition of human recombinant renin by the compounds of the invention

15 The enzymatic in vitro assay was performed in 384-well polypropylene plates (Nunc). The assay buffer consisted of 10 mM PBS (Gibco BRL) including 1 mM EDTA and 0.1% BSA. The incubates were composed of 50 μ L per well of an enzyme mix and 2.5 μ L of renin inhibitors in DMSO. The enzyme mix was premixed at 4°C and consists of the following components:

- 20 • human recombinant renin (0.16 ng/mL) • synthetic human angiotensin(1-14) (0.5 μ M)
• hydroxyquinoline sulfate (1 mM)


The mixtures were then incubated at 37°C for 3 h.

To determine the enzymatic activity and its inhibition, the accumulated Ang I was
25 detected by an enzyme immunoassay (EIA) in 384-well plates (Nunc). 5 μ L of the incubates or standards were transferred to immuno plates which were previously coated with a covalent complex of Ang I and bovine serum albumin (Ang I – BSA). 75 μ L of Ang I-antibodies in assay buffer above including 0.01% Tween 20 were added and a primary incubation made at 4 °C overnight. The plates were
30 washed 3 times with PBS including 0.01% Tween 20, and then incubated for 2 h at rt with an antirabbit-peroxidase coupled antibody (WA 934, Amersham). After washing the plates 3 times, the *peroxidase substrate* ABTS (2,2'-azino-di-(3-ethyl-

benzthiazolinsulfonate), was added and the plates incubated for 60 min at room temperature. After stopping the reaction with 0.1 M citric acid pH 4.3 the plate was evaluated in a microplate reader at 405 nm. The percentage of inhibition was calculated of each concentration point and the concentration of renin inhibition
5 was determined that inhibited the enzyme activity by 50% (IC_{50}). The IC_{50} -values of all compounds tested are below 10 μ M.

Claims

1. Compounds of the general formula I



wherein

Z, Y, X and W represent independently a nitrogen atom or a -CH- group; at least two of the Z, Y, X and W represent a -CH- group;

V represents a bond; $-(CH_2)_t-$; $-A-(CH_2)_s-$; $-CH_2-A-(CH_2)_t-$; $-(CH_2)_s-A-$; $-(CH_2)_2-A-(CH_2)_u-$; $-A-(CH_2)_v-B-$; $-CH_2-CH_2-CH_2-A-CH_2-$; $-A-CH_2-CH_2-B-CH_2-$; $-CH_2-A-CH_2-CH_2-B-$; $-CH_2-CH_2-CH_2-A-CH_2-CH_2-$; $-CH_2-CH_2-CH_2-CH_2-A-CH_2-$; $-A-CH_2-CH_2-B-CH_2-CH_2-$; $-CH_2-A-CH_2-CH_2-B-CH_2-$; $-CH_2-A-CH_2-CH_2-CH_2-B-$; or $-CH_2-CH_2-A-CH_2-CH_2-B-$;

A and B independently represent -O-; -S-; -SO-; -SO₂-;

U represents aryl; heteroaryl;

T represents $-CONR^1-$; $-(CH_2)_pOCO-$; $-(CH_2)_pN(R^1)CO-$; $-(CH_2)_pN(R^1)SO_2-$; or $-COO-$;

Q represents lower alkylene; lower alkenylene;

M represents hydrogen; cycloalkyl; aryl; heterocyclyl; heteroaryl;

L represents $-R^3$; $-\text{COR}^3$; $-\text{COOR}^3$; $-\text{CONR}^2\text{R}^3$; $-\text{SO}_2\text{R}^3$; $-\text{SO}_2\text{NR}^2\text{R}^3$;
 $-\text{COCH}(\text{Aryl})_2$;

5

R^1 represents hydrogen; lower alkyl; lower alkenyl; lower alkynyl; cycloalkyl;
aryl; cycloalkyl - lower alkyl;

R^2 and $R^{2'}$ independently represent hydrogen; lower alkyl; lower alkenyl;
10 cycloalkyl; cycloalkyl - lower alkyl;

R^3 represents hydrogen; lower alkyl; lower alkenyl; cycloalkyl; aryl; heteroaryl;
heterocyclyl; cycloalkyl - lower alkyl; aryl - lower alkyl; heteroaryl - lower alkyl;
heterocyclyl - lower alkyl; aryloxy - lower alkyl; heteroaryloxy - lower alkyl,
15 whereby these groups may be unsubstituted or mono-, di- or trisubstituted with
hydroxy, $-\text{OCOR}^2$, $-\text{COOR}^2$, lower alkoxy, cyano, $-\text{CONR}^2\text{R}^{2'}$, $-\text{CO-morpholin-4-yl}$,
 $-\text{CO-}((4\text{-loweralkyl})\text{piperazin-1-yl})$, $-\text{NH}(\text{NH})\text{NH}_2$, $-\text{NR}^4\text{R}^{4'}$ or lower alkyl,
with the proviso that a carbon atom is attached at the most to one heteroatom in
case this carbon atom is sp^3 -hybridized;

20

R^4 and $R^{4'}$ independently represent hydrogen; lower alkyl; cycloalkyl; cycloalkyl -
lower alkyl; hydroxy - lower alkyl; $-\text{COOR}^2$; $-\text{CONH}_2$;

k is the integer 0 or 1;

25

m and n represent the integer 0 or 1, with the proviso that in case m represents the
integer 1, n is the integer 0; in case n represents the integer 1, m is the integer 0; in
case k represents the integer 0, n represents the integer 0;

30 p is the integer 1, 2, 3 or 4;

r is the integer 1, 2, 3, 4, 5, or 6;

s is the integer 1, 2, 3, 4, or 5;

t is the integer 1, 2, 3, or 4;

u is the integer 1, 2, or 3;

v is the integer 2, 3, or 4;

5 and optically pure enantiomers, mixtures of enantiomers such as racemates, diastereomers, mixtures of diastereomers, diastereomeric racemates, mixtures of diastereomeric racemates, and the meso-form; as well as pharmaceutically acceptable salts, solvent complexes and morphological forms.

10 2. Compounds of general formula I according to claim 1 wherein Z, Y, X, W, V, U, T, Q, L, and M are as defined in general formula I and

k is 1

n is 0 and

15 m is 1

and optically pure enantiomers, mixtures of enantiomers such as racemates, diastereomers, mixtures of diastereomers, diastereomeric racemates, mixtures of diastereomeric racemates, and the meso-form; as well as pharmaceutically
20 acceptable salts, solvent complexes and morphological forms.

3. Compounds of general formula I according to claim 1 wherein Z, Y, X, W, V, U, T, Q, M, k, m, and n are as defined in general formula I and

25 L represents H; -COR^{3''}; -COOR^{3''}; -CONR^{2''}R^{3''};

R^{2''} and R^{3''} represent independently lower alkyl; lower cycloalkyl - lower alkyl, which lower alkyl and lower cycloalkyl-lower alkyl are undubstituted or mono-substituted with halogen, -CN, -OH, -OCOCH₃, -CONH₂, -COOH, or -NH₂, with
30 the proviso that a carbon atom is attached at the most to one heteroatom in case this carbon atom is sp³-hybridized,

and optically pure enantiomers, mixtures of enantiomers such as racemates, diastereomers, mixtures of diastereomers, diastereomeric racemates, mixtures of diastereomeric racemates, and the meso-form; as well as pharmaceutically acceptable salts, solvent complexes and morphological forms.

5

4. Compounds of general formula I according to claim 1 wherein Z, Y, X, W, V, U, L, k, m, and n are as defined in general formula I and

T represents -CONR¹-;

10

Q represents methylene;

M represents aryl or heteroaryl;

and optically pure enantiomers, mixtures of enantiomers such as racemates, diastereomers, mixtures of diastereomers, diastereomeric racemates, mixtures of
15 diastereomeric racemates, and the meso-form; as well as pharmaceutically acceptable salts, solvent complexes and morphological forms.

5. Compounds of general formula I according to claim 1 wherein V, U, T, Q, M, L, k, m, and n are as defined in general formula I and

20

Z, Y, X and W represent -CH-;

and optically pure enantiomers, mixtures of enantiomers such as racemates, diastereomers, mixtures of diastereomers, diastereomeric racemates, mixtures of
25 diastereomeric racemates, and the meso-form; as well as pharmaceutically acceptable salts, solvent complexes and morphological forms.

6. Compounds of general formula I according to claim 1 wherein Z, Y, X, W, V, Q, T, M, L, k, m, and n are as defined in general formula I and

30

U is a mono-, di-, or trisubstituted phenyl or heteroaryl, whereby the substituents are halogen, lower alkyl, lower alkoxy, CF₃

and optically pure enantiomers, mixtures of enantiomers such as racemates, diastereomers, mixtures of diastereomers, diastereomeric racemates, mixtures of diastereomeric racemates, and the meso-form; as well as pharmaceutically acceptable salts, solvent complexes and morphological forms.

5

7. The compounds according to any one of claims 1 to 6 selected from the group consisting of

(*rac.*)-(1*R**, 5*S**)-(3-acetyl-7-{3-[2-(2-bromo-5-fluorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

10

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-chlorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

15 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-*tert*-butylphenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,3,6-trifluorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

20

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,5-difluorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-[3-(2-*o*-tolylloxyethyl)phenyl]-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

25

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,3-dichlorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

30 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-chloro-5-methylphenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(3-chlorophenoxy)ethyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[3-(2-bromo-5-fluorophenoxy)propyl]phenyl}-
5 3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-chlorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

10 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-*tert*-butylphenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,3,6-trifluorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

15

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,5-difluorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-[3-(2-*o*-tolylloxypropyl)phenyl]-3,9-diazabicyclo-
20 [3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2,3-dichlorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

25 (*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(2-chloro-5-methylphenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(3-chlorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

30

(*rac.*)-(1*R**, 5*S**)-3-acetyl-7-{3-[2-(4-chlorophenoxy)propyl]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid methylphenethylamide;

(*rac.*)-(1*R**, 5*S**)-7-{3-[2-(2,6-dichloro-4-methylphenoxy)ethoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide;

5 (*rac.*)-(1*R**, 5*S**)-7-{3-[4-(2-fluoro-3-trifluoromethylphenoxy)butoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide;

(*rac.*)-(1*R**, 5*S**)-7-{3-[4-(2,6-dichloro-4-methylphenoxy)butoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide;

10

(*rac.*)-(1*R**, 5*S**)-7-{3-[4-(2-chloro-6-fluoro-3-methylphenoxy)butoxy]phenyl}-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxylic acid cyclopropyl-(3-methoxy-2-methylbenzyl)amide.

15

8. Pharmaceutical compositions containing at least one compound of any ones of claims 1 to 7 and usual carrier materials and adjuvants for the treatment or prophylaxis of disorders which are associated with a dysregulation of the renin-angiotensin system (RAS), comprising cardiovascular and renal diseases, hypertension, congestive heart failure, pulmonary hypertension, cardiac insufficiency, renal insufficiency, renal or myocardial ischemia, atherosclerosis, renal failure, erectile dysfunction, glomerulonephritis, renal colic, glaucoma, diabetic complications, complications after vascular or cardiac surgery, restenosis, complications of treatment with immunosuppressive agents after organ transplantation, and other diseases known to be related to the RAS.

20

25

9. A method for the treatment or prophylaxis of diseases which are related to the RAS comprising hypertension, congestive heart failure, pulmonary hypertension, cardiac insufficiency, renal insufficiency, renal or myocardial ischemia, atherosclerosis, renal failure, erectile dysfunction, glomerulonephritis, renal colic, glaucoma, diabetic complications, complications after vascular or cardiac surgery,

30

restenosis, complications of treatment with immunosuppressive agents after organ transplantation, and other diseases which are related to the RAS, which method comprises administering a compound according to any one of claims 1 to 7 to a human being or animal.

5

10. The use of compounds according to any one of claims 1 to 7 for the treatment or prophylaxis of diseases which are associated with the RAS comprising hypertension, congestive heart failure, pulmonary hypertension, cardiac insufficiency, renal insufficiency, renal or myocardial ischemia, atherosclerosis, renal failure, erectile dysfunction, glomerulonephritis, renal colic, glaucoma, diabetic complications, complications after vascular or cardiac surgery, restenosis, complications of treatment with immunosuppressive agents after organ transplantation, and other diseases known to be related to the RAS.

11. The use of one or more compounds of any one of claims 1 to 7 in combination with other pharmacologically active compounds comprising ACE inhibitors, angiotensin II receptor antagonists, endothelin receptor antagonists, vasodilators, calcium antagonists, potassium activators, diuretics, sympatholitics, beta-adrenergic antagonists, alpha-adrenergic antagonists, and neutral endopeptidase inhibitors, for the treatment of disorders as set forth in any one of claims 9 to 11.

INTERNATIONAL SEARCH REPORT

International Application No

EP2004/004373

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D471/08 A61K31/4995 A61P13/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97/09311 A (HOFFMANN LA ROCHE) 13 March 1997 (1997-03-13) cited in the application page 64, line 12 - page 73, line 30; claim 1	1-11
P,X	----- WO 03/093267 A (REMEN LUBOS ; WELLER THOMAS (CH); BUR DANIEL (CH); FISCHLI WALTER (CH)) 13 November 2003 (2003-11-13) page 476, line 4 - line 32; claim 1; examples 1-743 -----	1-11



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

- *T* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the International search

26 July 2004

Date of mailing of the International search report

03/08/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Seelmann, I

INTERNATIONAL SEARCH REPORT

Information on patent family members

Original Application No

EP2004/004373

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9709311	A	13-03-1997	AT 242213 T 15-06-2003
		AU 708616 B2 05-08-1999	
		AU 6743296 A 27-03-1997	
		BR 9610385 A 06-07-1999	
		CA 2230931 A1 13-03-1997	
		CN 1202152 A 16-12-1998	
		CZ 9800684 A3 14-10-1998	
		DE 59610509 D1 10-07-2003	
		DK 863875 T3 01-12-2003	
		WO 9709311 A1 13-03-1997	
		EP 0863875 A1 16-09-1998	
		ES 2201192 T3 16-03-2004	
		HU 9900926 A2 28-09-1999	
		IL 123293 A 24-06-2003	
		JP 11500447 T 12-01-1999	
		MA 23967 A1 01-04-1997	
		NO 980954 A 28-04-1998	
		NZ 315677 A 28-02-2000	
		PL 325425 A1 20-07-1998	
		PT 863875 T 31-10-2003	
		RU 2167865 C2 27-05-2001	
		TR 9800409 T1 21-05-1998	
		TW 474932 B 01-02-2002	
		US 6051712 A 18-04-2000	
		US 6150526 A 21-11-2000	
		ZA 9607424 A 07-03-1997	
WO 03093267	A	13-11-2003	WO 03093267 A1 13-11-2003